

# The Role of Resveratrol as Potent Antioxidant for Anti-Aging Formulation Development: A Review

*by* Widji Soeratri

---

**Submission date:** 07-Jan-2020 02:29PM (UTC+0800)

**Submission ID:** 1239715083

**File name:** DIJR-DEC-007\_1.pdf (497.04K)

**Word count:** 2904

**Character count:** 16697

## The Role of Resveratrol as Potent Antioxidant for Anti-Aging Formulation Development: A Review

Dita Nurlita Rakhma<sup>\*1</sup>, Noorma Rosita<sup>2</sup>, Widji Soeratri<sup>3</sup>

<sup>1,2,3</sup>Department of Pharmaceutics, Faculty of Pharmacy, Airlangga University, Indonesia  
60286, Surabaya, Indonesia

\*Corresponding Author E-mail: [ditanurlitar@gmail.com](mailto:ditanurlitar@gmail.com)

### Abstract

The occurrence of skin aging process is not the same in each individuals. The changes in skin appearance can affect in self-confidence, especially for those who experience the aging process faster than their age called premature aging. Therefore it is necessary therapy to prevent, repair and maintain the structure and function of aging skin. Resveratrol is a potent antioxidant that has activity as skin anti-aging. The limitations of the physicochemical properties of Resveratrol become the focus of attention for formulation of anti-aging development.

**Keywords :** Anti-aging, Antioxidant, Resveratrol, Lipid Nanoparticles, Vesicles, Nanocarriers

### I. INTRODUCTION

Skin aging is a complex biological process that will be experienced in every human (1). The aging process is a gradual process of slowing body's tissue ability to repair or maintain its structure and normal functioning. Skin is the most outer layer of the body which several important roles to protect humans body (2). In certain people aging process can occur according to the ages, but there are some people who experience skin aging process earlier than ages (the process of premature aging) and some are experiencing the aging process more slowly than the

25s. This shows that aging process in each individuals depends on many factors affect. Extracellular matrix (ECM) plays an important role in maintaining skin integrity. ECM has three main components: elastic fibers, collagen and proteoglycans (3). When the aging factor appears, the extracellular matrix will be remodeling (4). The visible changes in skin aging such as dry, coarse, wrinkles accompanied by lines of facial expression can affect appearance showing that the person has entered old age. These changes eventually decline self-confidence, especially in those who experience the aging process faster than their age (5). Therefore, it requires management of both invasive and non-invasive therapy to prevent, improve or maintain the structure and function of aging skin.

### II. SKIN AGING

Aging process can be classified by intrinsic factors (chronological aging) and extrinsic factors (premature aging).

#### 1. Intrinsic aging

This process also called chronological aging, a natural aging process caused by various physiological factors of the body itself, such as genetic, hormonal, and ethnic. In hormonal factors, estrogen has an influence in the synthesis of collagen by fibroblasts, which then stimulate synthesis of hyaluronic acid and increase extracellular matrix. In menopause, there is a condition of hypoenestrogenism that affects the elasticity of the skin. While in the ethnic factor, the aging process comes with pigmentation. High levels of melanin pigmentation will protect from the cumulative effect of photoaging. Black ethnic has a greater number of lipids and is compact and so resistant to photoaging. Likewise, the Asian has a slower skin aging ability than the Caucasian. Telomere, a small DNA sequence located at the end of a chromosome is considered an essential element in the intrinsic aging process. Along with the aging process, continuous replication will lead to the shortening of DNA structure that can be repaired by telomerase. Telomere maintenance by telomerase action will inhibit the aging process, but this action can lead to carcinogenesis (6).

#### 2. Extrinsic aging

The aging process that occurs due to various factors from the environment is called premature aging. It is caused by several factors such as ultraviolet radiation (UV), air humidity, temperature, pollutant. A person can experience aging faster than its age due to the ongoing exposure of these environmental factors. Free radicals factor obtained from UV exposure is a major factor that accelerates the aging process called photoaging. Free radicals can cause damage to skin such as producing Reactive Oxygen Species (ROS) that can cause DNA damage, induce the enzyme matrix metalloproteinase (MMP) that leads to collagen degradation (7).

#### A. Treatment of Skin Aging


To overcome the problem of skin aging, begins with the prevention of UV exposure. This can be done by using sunblock or sunscreen with chemical or physical filters. The next step that can be done if signs of premature aging have visible:

1. Use active ingredients to reduce signs of premature aging of the skin.
2. If the condition of premature aging on the skin more serious invasive therapy required.

3. To improve the condition of the skin it can use cosmetics such as antioxidants, retinoids, and alpha hydroxy acids that are used routinely to show an improvement in skin conditions (8).

### 8. Resveratrol as Skin Anti-Aging

Resveratrol (3) a potent antioxidant that is currently being investigated for a variety of disease conditions. Resveratrol (3,5,4'-trihydroxystilbene) (Fig.1) is a naturally occurring phytoalexin antioxidant present in grapes, berries, peanuts and red wine (9). In 1940, resveratrol was first derived from the white hellebore root (*Veratrum grandiflorum* O. Loes) and nearly 75 years later, it was found in the roots of *Polygonum cuspidatum* (10). This plant has been used for traditional medicine where it is valued for diverse therapeutic effects (11).



11 Figure 1. Structures of trans-resveratrol (trans-R) and cis-resveratrol (cis-) (12)

Resveratrol characterized by solid off-white powder with molecular formula  $C_{14}H_{12}O_3$ , molecular weight (8) of 228.25 g.mol<sup>-1</sup>, melting point between 253-255 °C, and a fat soluble-compound. Resveratrol consists of two structural isomers: cis- (Z) and trans- (E) (Fig. 1). The trans-resveratrol biologically more active than the Cis-resveratrol (73). Exposure to the sunlight, heat or UV radiation, can turn trans-resveratrol into isomerisation cis form (14). Resveratrol exhibits high membrane permeability and can be considered a class-II compound in the Biopharmaceutical Classification System (15).

Resveratrol effectively inhibited AP-1 and NF-κB expression (16). AP-1 is a main transcription factor (1) responsible for the production of metalloproteinases (MMP) enzymes that contribute (2) to collagen degradation. Increased (18) collagen degradation is a major cause of wrinkles. While NF-κB produces a pro-inflammatory mediator that also plays a role in skin aging. Other studies have shown that resveratrol prevents mitochondrial dysfunction (17). The elevation of mitochondrial function decreases the regeneration of body cells that can accelerate aging (18), and thus application of resveratrol becomes the primary key as anti-aging.

22 *In vivo* studies have shown that topical use of resveratrol in hairless mice reduces skin inflammation, inhibits cyclooxygenase (COX) ornithine decarboxylase (ODC) induction, and formation of hydrogen peroxide due to UV-B exposure (19).

Based on its chemical structure, resveratrol has a stilben group similar to estrogen diethylstilbesterol. Therefore, resveratrol also has activity as phytoestrogens and estrogen beta receptor agonist (ERβ). In postmenopausal women, clinical signs of skin aging will show (1) due to the loss of estrogen production. The use of resveratrol as phytoestrogens becomes potential to overcome the clinical signs of skin aging in the absence of side risks (20).

### C. Topical Formulation of Resveratrol as Anti-Aging

Resveratrol is a polyphenol compound with many several health benefits. Resveratrol has many mechanisms as a powerful antioxidant to improve the functioning of organs systems including the skin. Resveratrol plays an important role as a topical ingredient for treating the skin as it can prevent and correct the clinical (4) signs of aging (21). However, further innovation in the delivery system is needed to overcome the limitations of resveratrol in order (10) to maintain the stability and effectiveness of resveratrol into the skin (22).

#### 1. Lipid nanoparticles

Solid Lipid (9) nanoparticles (SLN) is the first generation of lipid nanoparticles that has developed since 1990 (23). SLN is produced by replacing the liquid lipid component (oil) from oil / water emulsions with solid lipids (14). SLN has ability to protect unstable chemical ingredients and prolonging drug release (25). Nanostructured Lipid Carriers (NLC) is the latest generation of Solid Lipid Nanoparticle (Fig.2). NLC was developed to overcome the weakness of SLN due to its limited loading and the occurrence of drug expulsion during storage. NLC consists of a certain amount of lipid matrix and liquid lipids. The NLC remains in solid formation by controlling the oil content added to the formulation, so that the controlled drug release drug for NLC can be achieved.




Figure 2. Structure of SLN and NLC (26)

In 2012, Gokce et al (27) conducted a study to determine the effectiveness of resveratrol formulated in SLN and NLC systems with Compritol as solid lipids and miglyol as lipid liquids. The results showed that the use of topical resveratrol in SLN and NLC systems increasing resveratrol penetration into the dermis and protect fibroblasts from exposure to ROS.

## 2. Vesicles

Vesicular carriers can be divided into two main groups, namely liposomes and niosomes (28). Liposomes are vesicular structures consist of a hydrophobic lipid bilayer membrane surrounding an aqueous core. Hydrophobic lipid bilayer membrane composed of a phospholipid and cholesterol (29). Resveratrol loaded by liposomes achieve incorporation percentage higher than 70%, low zeta potential during storage, and increase cell proliferation under the stress condition induced by UV-B light (30). These result showing good capability in incorporation stability and photoprotective effect.

Niosomes are non-ionic surfactants vesicle in aqueous media resulting in closed bilayer structures. Niosomes have more advantages over liposomes because of higher chemical stability and penetration (31). An ex vivo studies conducted by Negi et al in 2017 (32), show high permeation and resveratrol deposition in skin when compared with ordinary resveratrol. In addition, the study also showed that resveratrol entrapped niosomal gel significantly reduced edema in the rat paw induced carragenan model.

## 3. Protein based nanocarriers

Protein-based nanocarriers are being developed due to biodegradability, biocompatibility, high nutritional value, low cytotoxicity, abundant renewable sources and extraordinary binding capacity of various drugs (33). Silk is a protein polymer produced by insects and spiders. Silk protein has several advantages for drug delivery due to its biocompatibility, slow biodegradability, controllable structure and morphology and less inflammatory (35). Silk protein from silkworms deinking consists of two proteins, fibroin and sericin. (36). Fibroin is the core of silk of a fibrous protein, while sericin is a protein that wrap fibroin fibers with a successive sticky layers to help the formation of cocoons (37).

An in vitro cytotoxic study has been conducted by Sukhtam in 2017, about resveratrol loaded sericin carriers, it is known that there is no cytotoxic effect against normal skin fibroblast cells. This result suggests that resveratrol load sericin is safely applied to the manufacture of therapeutics and cosmetics.

## III. CONCLUSION

Based on several studies, it is known that resveratrol can be formulated in various delivery systems to maintain its stability, safety and effectivity as anti-aging. These factors are important in the formulation development of drugs and cosmetics.

## REFERENCES

1. Papakonstantinou E, Roth M, Karakiulakis G. Hyaluronic acid: A key molecule in skin aging. *Dermato-endocrinology*, 2012; 4(3): 253-8.
2. Tobin, DJ. Introduction to skin aging. *Journal of tissue viability*, 2017; 26(1): 37-46.
3. Bradley EJ, Griffiths C, Sherratt MJ, Bell M, Watson REB. Over-the-counter anti-ageing topical agents and their ability to protect and repair photoaged skin. *Maturitas*, 2015; 80: 265-72.
4. Rinnerthaler M, Bischof J, Streubel MK, Trost A, Richter K. Oxidative Stress in Aging Human Skin. *Biomolecules*, 2015; 5: 545-89
5. Farage MA, Miller KW, Elsner P, Maibach HI. Characteristics of the aging skin. *Advances in wound care*, 2013; 2(1): 5-10.
6. Silva MR, Celeim LR, Silva SR, Costa APF. Anti-aging cosmetics: Facts and controversies. *Clinics in Dermatology*, 2013; 31: 750-8

7. Berneburg M, Plettenberg H, Krutmann J. Photoaging of human skin. *Photodermatol Photoimmunol Photomed* 2000; 16: 239-44
8. Silva MR, Celem LR, Silva SR, Costa APF. Anti-aging cosmetics: Facts and controversies. *Clinics in Dermatology*. 2013; 31: 750-8
9. Ndiaye M, Philippe C, Mukhtar H, Ahmad, N. The grape antioxidant resveratrol for skin disorders: promise, prospects, and challenges. *Archives of biochemistry and biophysics*, 2011; 508(2): 164-70.
10. Baur JA, Sinclair, DA. Therapeutic potential of resveratrol: the in vivo evidence. *Drug Discovery*, 2006; 5:493-506
11. Farris P, Krutman J, Li YH, , McDaniel, Krolj, Y. Resveratrol: a unique antioxidant offering a multi-mechanistic approach for treating aging skin. *J Drugs Dermatol*, 2013; 12(12): 1389-94
12. Anisimova, NY, Kiselevsky, M.V, Sosnov, A V, Sadovnikov, SV, Stankov, IN, Gakh, AA. Trans-, cis-, and dihydro-resveratrol: a comparative study. *Chemistry Central Journal*., 2011; 5(1): 1-6.
13. Amri A, Chaumeil JC, Sfar S, Charrueau. Administration of resveratrol: What formulation solutions to bioavailability limitations?. *Journal of Controlled Release*, 2012; 158: 182-93
14. Davidov-Pardo G, McClements DJ. Resveratrol Encapsulation Designing Delivery Systems To Overcome Solubility, Stability and Bioavailability Issues. *Trends in Food Science & Technology*, 2014; 38: 88-103
15. Amidon GL, Lennernäs H, Shah VP, Crison JR. A theoretical basis for a biopharmaceutic drug classification: the correlation of in vitro drug product dissolution and in vivo bioavailability. *Pharm. Res*, 1995; 12: 413-20.
16. Yang T, Wang L, Zhu M, Zhang L, Yan L. Properties and molecular mechanisms of resveratrol: a review. *Pharmazie*, 2015; 70: 501-6
17. Farris P, Krutman J, Li YH, , McDaniel, Krolj, Y. Resveratrol: a unique antioxidant offering a multi-mechanistic approach for treating aging skin. *J Drugs Dermatol*, 2013; 12(12): 1389-94
18. Krutmann J, Schroeder P. Role of mitochondria in photoaging of human skin: the defective powerhouse model. In *Journal of Investigative Dermatology Symposium Proceedings*, 2009; 14(1): 44-9. Elsevier.
19. Ndiaye M, Philippe C, Mukhtar H, Ahmad, N. The grape antioxidant resveratrol for skin disorders: promise, prospects, and challenges. *Archives of biochemistry and biophysics*, 2011; 508(2): 164-70.
20. Farris P, Krutman J, Li YH, , McDaniel, Krolj, Y. Resveratrol: a unique antioxidant offering a multi-mechanistic approach for treating aging skin. *J Drugs Dermatol*, 2013; 12(12): 1389-94
21. Sun R, Zhao G, Ni S, Xia Q. Lipid based nanocarriers with different lipid compositions for topical delivery of resveratrol: comparative analysis of characteristics and performance, 2014; 24(6): 591-600
22. Farris P, Krutman J, Li YH, , McDaniel, Krolj, Y. Resveratrol: a unique antioxidant offering a multi-mechanistic approach for treating aging skin. *J Drugs Dermatol*, 2013; 12(12): 1389-94
23. Pardeike J, Hommoss A, Müller RH. Lipid nanoparticles (SLN, NLC) in cosmetic and pharmaceutical dermal products. *International Journal of Pharmaceutics*, 2009; 366: 170-84
24. Pardeike J, Schwabe K, Müller RH. Influence of nanostructured lipid carriers (NLC) on the physical properties of the Cutanova Nanorepair Q10 cream and the in vivo skin hydration effect. *International Journal of Pharmaceutics*, 2010; 396: 166-73
25. Huang ZR, Lin YK, Fang JY. Biological and pharmacological activities of squalene and related compounds: potential uses in cosmetic dermatology. *Molecules*, 2009; 14(1): 540-554.
26. Muller RH, Shegokar R, M Keck, 20 years of lipid nanoparticles (SLN & NLC): present state of development & industrial applications. *Current drug discovery technologies*, 2011; 8(3): 207-227.
27. Gokce EH, Korkmaz E, Dellera E, Sandri G, Bonferoni MC, Ozer O. Resveratrol-loaded solid lipid nanoparticles versus nanostructured lipid carriers: evaluation of antioxidant potential for dermal applications. *International journal of nanomedicine*, 2012 ;7: 1841-50.
28. Sinico, C., & Fadda, A. M. (2009). Vesicular carriers for dermal drug delivery. *Expert opinion on drug delivery*, 6(8), 813-825
29. Arora, N, Agarwal, S, Murthy, RSR. Latest technology advances in cosmeceuticals. *Int J Pharm Sci Drug Res*, 2012; 4(3), 168-82
30. Caddeo, C, Teskač, K, Sinico, C, Kristl. Effect of resveratrol incorporated in liposomes on proliferation and UV-B protection of cells. *International journal of pharmaceutics*, 2008; 363(1): 183-91.
31. Pando D, Matos M, Gutiérrez G, Pazos C. (2015). Formulation of resveratrol entrapped niosomes for topical use. *Colloids and Surfaces B: Biointerfaces*, 2015; 128: 398-404.
32. Negi P, Aggarwal M, Sharma G, Rathore C, Sharma G, Singh B, Katare OP. Niosome-based hydrogel of resveratrol for topical applications: An effective therapy for pain related disorder (s). *Biomedicine & Pharmacotherapy*, 2017; 88: 480-487.
33. Suktham, K, Koobkobkruad, T, Wutikhun, T, Surassmo, S. Efficiency of resveratrol-loaded sericin nanoparticles: Promising bionanocarriers for drug delivery. *International Journal of Pharmaceutics*, 2017, 1-6.



34. Elzoghby, AO, Samy, WM, Elgindy, NA. Protein-based nanocarriers as promising drug and gene delivery systems. *Journal of controlled release*, 2012; 161(1): 38-49
35. Numata, K, Kaplan, DL. Silk-based delivery systems of bioactive molecules. *Advanced drug delivery reviews*, 2010; 62(15), 1497-508.
36. Suktham, K, Koobkobkruad, T, Wutikhun, T, Surassmo, S. Efficiency of resveratrol-loaded sericin nanoparticles: Promising bionanocarriers for drug delivery. *International Journal of Pharmaceutics*, 2017, 1-6.
37. Altman GH, Diaz F, Jakuba C, Calabro T, Horan, RL, Chen J, Lu H, Richmond J, Kaplan DL. Silk-based biomaterials. *Biomaterials*, 2003;24(3): 401-6

# The Role of Resveratrol as Potent Antioxidant for Anti-Aging Formulation Development: A Review

## ORIGINALITY REPORT

25%

SIMILARITY INDEX

13%

INTERNET SOURCES

22%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.dermagist.com](http://www.dermagist.com)

Internet Source

4%

2

"Textbook of Aging Skin", Springer Nature, 2017

Publication

3%

3

Mary Ndiaye, Carol Philippe, Hasan Mukhtar, Nihal Ahmad. "The grape antioxidant resveratrol for skin disorders: Promise, prospects, and challenges", Archives of Biochemistry and Biophysics, 2011

Publication

2%

4

A. Amri, J.C. Chaumeil, S. Sfar, C. Charrueau. "Administration of resveratrol: What formulation solutions to bioavailability limitations?", Journal of Controlled Release, 2012

Publication

2%

5

Ahmed O. Elzoghby, Mayada M. Elgohary, Nayra M. Kamel. "Implications of Protein- and Peptide-Based Nanoparticles as Potential Vehicles for Anticancer Drugs", Elsevier BV,

2%

6

Daniel Pando, María Matos, Gemma Gutiérrez, Carmen Pazos. "Formulation of resveratrol entrapped niosomes for topical use", *Colloids and Surfaces B: Biointerfaces*, 2015

Publication

---

7

Amri, A.. "Administration of resveratrol: What formulation solutions to bioavailability limitations?", *Journal of Controlled Release*, 20120310

Publication

---

8

Valentina Venuti, Carmela Cannavà, Maria Chiara Cristiano, Massimo Fresta et al. "A characterization study of resveratrol/sulfobutyl ether- $\beta$ -cyclodextrin inclusion complex and in vitro anticancer activity", *Colloids and Surfaces B: Biointerfaces*, 2014

Publication

---

9

Gökçe, Evren H., Evren Algın Yapar, Sakine Tuncay Tanrıverdi, and Zengin Zeynep. "Nanocarriers in cosmetology", *Nanobiomaterials in Galenic Formulations and Cosmetics*, 2016.

Publication

---

10

[estudogeral.sib.uc.pt](http://estudogeral.sib.uc.pt)

Internet Source

---

1%

1%

1%

1%

1%



11	ccj.springeropen.com Internet Source	1 %
12	Annish Jain, Sumit K. Singh, Shailendra K. Arya, Subhas C. Kundu, Sonia Kapoor. "Protein Nanoparticles: Promising Platforms for Drug Delivery Applications", ACS Biomaterials Science & Engineering, 2018 Publication	1 %
13	Kai Su, Chunming Wang. "Recent advances in the use of gelatin in biomedical research", Biotechnology Letters, 2015 Publication	1 %
14	Percutaneous Penetration Enhancers Chemical Methods in Penetration Enhancement, 2016. Publication	1 %
15	Yen-Chih Lin. "Spatially Controlled Delivery of Neurotrophic Factors in Silk Fibroin–Based Nerve Conduits for Peripheral Nerve Repair :", Annals of Plastic Surgery, 08/2011 Publication	<1 %
16	www.em-consulte.com Internet Source	<1 %
17	www.tandfonline.com Internet Source	<1 %
18	Bioactive Dietary Factors and Plant Extracts in Dermatology, 2013.	<1 %

---

19 Eunson Hwang, Sang-Yong Park, Hyun Ji Lee, Tae Youp Lee, Zheng-wang Sun, Tae Hoo Yi. "Gallic Acid Regulates Skin Photoaging in UVB-exposed Fibroblast and Hairless Mice", Phytotherapy Research, 2014

Publication

---

20 John M. Pezzuto. "Resveratrol as an Inhibitor of Carcinogenesis", Pharmaceutical Biology, 2008

Publication

---

21 [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

---

22 [vdocuments.site](http://vdocuments.site)

Internet Source

---

23 [e-sciencecentral.org](http://e-sciencecentral.org)

Internet Source

---

24 Kim, Min Kyoung, Chae Young Bang, Gwang Jun Yun, Hyang-Yu Kim, Young Pyo Jang, and Se Young Choung. "Anti-wrinkle effects of Seungma-Galgeun-Tang as evidenced by the inhibition of matrix metalloproteinase-1 production and the promotion of type-1 procollagen synthesis", BMC Complementary and Alternative Medicine, 2016.

Publication

---

25 Andrea S. Gobin, Victoria E. Froude, Anshu B.

Mathur. "Structural and mechanical characteristics of silk fibroin and chitosan blend scaffolds for tissue regeneration", Journal of Biomedical Materials Research Part A, 2005

Publication

<1%

26

Vierkötter, Andrea, and Jean Krutmann. "Environmental influences on skin aging and ethnic-specific manifestations", Dermato-Endocrinology, 2012.

Publication

<1%

27

Poonam Negi, Madhur Aggarwal, Gajanand Sharma, Charul Rathore, Gaurav Sharma, Bhupinder Singh, O.P. Katare. "Niosome-based hydrogel of resveratrol for topical applications: An effective therapy for pain related disorder(s)", Biomedicine & Pharmacotherapy, 2017

Publication

<1%

28

"Cosmetic Dermatology", Wiley, 2010

Publication

<1%

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

# The Role of Resveratrol as Potent Antioxidant for Anti-Aging Formulation Development: A Review

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5